

AD-A080 181

TEXAS A AND M UNIV COLLEGE STATION DEPT OF CHEMISTRY

F/8 7/4

ELECTRON TRANSFER REACTIVITY ON NOVEL DIRHENIUM AND DIRHODIUM T--ETC(U)

JAN 80 M TSUTSUI, R BOBSEIN, K M KADISH

N00014-75-C-0417

UNCLASSIFIED

TR-32

NL

1 OF 1
AD
A080181




END

DATE
FILMED

3 - 80

DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

DA080181

DDC FILE COPY

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 32	2. GOVT ACCESSION NO. 85	3. RECIPIENT'S CATALOG NUMBER SECRET
4. TITLE (and Subtitle) (6) Electron Transfer Reactivity of Novel Dirhenium and Dirhodium Tetraazaannulene Complexes.	5. TYPE OF REPORT & PERIOD COVERED (9) Interim rept.	
6. PERFORMING ORG. REPORT NUMBER		7. AUTHOR(s) (10) M./Tsutsui, Rex/Bobsein, K.M./Kadish, L.A./Bottomley D./Schaeper
8. CONTRACT OR GRANT NUMBER(s) (15) No 014-75-C-0417		9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Texas A&M University College Station, Texas 77843
10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-559		11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217 (14) TR-32
12. REPORT DATE (11) Jan 1980		13. NUMBER OF PAGES 13
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (12/15)		15. SECURITY CLASS. (of this report) 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release; Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) DDC RECEIVED FEB 5 1980 RECEIVED A		
18. SUPPLEMENTARY NOTES Submitted for publication in Inorganic Chimica Acta		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Electron Transfer Reactivities, 5,7,12,14-tetramethyldibenzo[b,i][1,4,8,11]tetraaza[14]annulene, [1,4,8,11]tetraaza[14]annulene, tetraphenylporphyrin, [Re(CO) ₃] ₂ TMTAA, [Rh(CO) ₂] ₂ TMTAA, [Re(CO) ₃] ₂ TAA, (CO) ₃ TPP		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The electron transfer reactivities of dirhenium and dirhodium carbonyl complexes of the macrocyclic ligands, 5,7,12,14-tetramethyldibenzo[b,i][1,4,8,11]tetraaza[14]annulene, (TMTAA), dibenzo [b, i][1,4,8,11]tetraaza[14]annulene, (TAA), and tetraphenylporphyrin, (TPP) were investigated by cyclic voltammetry and differential pulse polarography. Potentials are reported for the electrochemical generation of \dot{R}^+ cation and \dot{R}^- anion radicals of these complexes in the solvents dichloromethane and dimethylformamide. The potential differences between the \dot{R}^+ radical reactions of the metallomacrocyclic complexes were compared to the po-		

DD FORM 1473
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

407 997
80 2 4 099

tential differences of the π radical reactions of the free macrocycles.

Abstract

The electron transfer reactivities of dirhenium and dirhodium carbonyl complexes of the macrocyclic ligands 5,7,12,14-tetramethyl dibenzo[b,i] [1,3,8,11]tetraaza[14]annulene, (TMTAA), dibenzo[b,i] [1,3,8,11]tetraaza [14]annulene, (TAA), and tetraphenylporphyrin, (TPP) were investigated by cyclic voltammetry and differential pulse polarography. Potentials are reported for the electrochemical generation of π cation and π anion radicals of these complexes in the solvents dichloromethane and dimethylformamide. The potential differences between the π radical reactions of the metallo macrocyclic complexes were compared to the potential differences of the π radical reactions of the free macrocycles.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Availand/or special
A	

The synthesis and structural characterization of dirhodium¹ and dirhenium² carbonyl complexes of the macrocyclic ligand 5,7,12,14-tetramethyl dibenzo [b,i] [1,4,8,11]tetraaza[14]annulene, (TMTAA) and dirhenium carbonyl complexes³ of dibenzo [b,i] [1,4,8,11] tetraaza[14]annulene, (TAA) have recently been completed. These compounds are of interest both in terms of their structural similarity to the natural porphyrins⁴ and their potential role in catalysis.⁵

Complexes of H₂TAA and H₂TMTAA with a wide variety of +2 and +3 metal ions have been synthesized.⁶⁻⁷ The metal(II) or metal(III) ion replaces the two hydrogens from H₂TAA or H₂TMTAA and forms tetradentate N₄-coordinated metal complexes in a manner similar to metalloporphyrins. Resonance Raman studies of M(II)TMTAA (M = Mn, Fe, Co, Ni, Cu or Zn) reveal many similarities between the macrocyclic complexes of the TMTAA ligand and those of the metalloporphyrins.⁸ Similarly, electrochemical oxidation of NiTMTAA⁹ or [Mn(III)TMTAA]⁺¹⁰ yields π cation radicals similar to those observed for metalloporphyrin oxidation.¹¹ However, to date, all comparisons between the porphyrins and synthetic macrocycles have involved monometallic M(II) and M(III) complexes in which the metal is complexed to all four donor nitrogen atoms. In this study we have investigated the redox properties of four different dimetallic M(I) macrocyclic complexes in which each metal atom is bonded to either two or three nitrogen donor atoms. We have used cyclic voltammetry to probe energy levels of the macrocyclic complexes [Re(CO)₃]₂TMTAA and compared their half wave potentials to those obtained for the oxidation-reduction of [Re(CO)₃]₂TAA and the dirhenium complex of tetraphenylporphyrin, [Re(CO)₃]₂TPP. The four complexes differ significantly not only in terms of the number of coordinated nitrogens but also in terms of symmetry and degree of metal-metal interactions. (See Figure 1)

Experimental

Chemicals: H_2TAA , $[\text{Re}(\text{CO})_3]_2\text{TAA}$, $[\text{Re}(\text{CO})_2]_2\text{TMTAA}$ and $[\text{Rh}(\text{CO})_2]_2\text{TMTAA}$ were prepared by literature methods.¹⁻³

Dichloromethane (CH_2Cl_2) (Fisher, distilled from P_2O_5) and N,N-dimethylformamide (DMF) (Fisher reagent) were used as solvents for electrochemical studies. The supporting electrolyte, tetrabutylammonium fluoroborate (TBAT) (Fluka), was recrystallized from methanol and dried under reduced pressure prior to use.

Cyclic Voltammetry. Cyclic voltammograms were obtained either with a PAR 174 polarographic analyzer, in conjunction with a Houston Instruments 2000 X-Y recorder, or a PAR 173 potentiostat and 175 Universal Programmer on an oscilloscope. A three-electrode system was used, consisting of platinum working and counter electrodes and a commercial saturated lithium calomel electrode (SLCE) that was separated from the bulk of the solution by a bridge filled with solvent and supporting electrolyte. Solutions in the bridge were changed periodically to avoid aqueous contamination from entering the cell via the SLCE. All solutions were purged of oxygen by passing purified nitrogen through them for 10 min immediately prior to running cyclic voltammograms. For dichloromethane solutions, the nitrogen was solvent saturated prior to entering the cell. After degassing, a blanket of nitrogen was maintained over the solution.

For reversible electrode reactions, the $E_{1/2}$ values were measured as that potential lying midway between the oxidation and reduction peak for a given couple. For several oxidations in DMF, no reverse cathodic peak was obtained. In these instances the potentials reported are the anodic peak potentials, E_p , recorded at 1V/sec. These potentials were scan rate dependent and shifted anodically by approximately $30/n$ mV for each tenfold increase in scan rate.

Results and Discussion

Figure 2 illustrates a typical cyclic voltammogram of H_2TAA in DMF. At fast

sweep rates, the half wave potentials were independent of scan rate and $i_p/v^{1/2}$ was constant as expected for a diffusion controlled oxidation-reduction. At slow sweep rates, no reverse peak was observed and E_p shifted along the potential axis by 30-40mV per tenfold increase in scan rate. These criteria are indications of an electron transfer followed by a chemical reaction (EC).¹² This mechanism was further confirmed by the invariant peak potential shapes of $E_p - E_{p/2} = 60 \pm 5$ mV (Fig 2). Peak heights were approximately equal for the oxidation and the reduction processes, confirming the hypothesis that equal numbers of electrons were transferred in each reaction.

The rate constant for the following chemical reaction was largest in the absence of a metal ion. For the slow chemical reactions, reversible cyclic voltammograms could be obtained at rapid scan rates. However, in both CH_2Cl_2 and DMF the followup chemical reactions of $[\text{H}_2\text{TAA}]^+$ and $[\text{H}_2\text{TMTAA}]^+$ were so rapid that even at scan rates of 100V/sec, no reverse reduction peaks were obtained. No attempt was made to ascertain the nature of the chemical reaction.

Potentials for the electrode reactions of each ligand in DMF and its corresponding dirhenium or dirhodium carbonyl complexes are listed in Table I, while Table II lists the reversible potentials of the dirhodium and dirhenium complexes in CH_2Cl_2 . Changing the ligand from H_2TPP to H_2TAA has little effect on the oxidation process to yield cation radicals. In contrast, formation of the anion radical is favored by 0.68V when the ligand is H_2TPP compared to H_2TAA and by over 1.5 V when compared to H_2TMTAA . A similar shift in reduction potential is observed on going from $[\text{Re}(\text{CO})_3]_2\text{TPP}$ ($E_{1/2} = -0.66\text{V}$) to $[\text{Rh}(\text{CO})_3]_2\text{TMTAA}$ ($E_{1/2} > -2.20\text{V}$).

Most metal complexes with H_2TMTAA are "Saddle shaped"^{4,8,13} (The only known exception is the palladium complex of the H_2TMTAA ligand.¹⁴) with the o-phenylene rings and diiminate framework displaced in opposite directions from the plane of the four nitrogens. This saddle shape configuration allows

the two Re(I) or Rh(I) atoms to sit atop the complex with both atoms in close proximity. The dimetallic complexes of TMTAA are thus symmetrical molecules where the metal atoms are each bound to two of the four adjacent donor nitrogen atoms. The metal ions reside in approximately the same plane as the pentane-diiminato chelate ring containing the two coordinated nitrogen atoms. Structural characterization of these complexes show a Rh-Rh separation of 3.086 \AA ¹ and a Re-Re separation of 4.11 \AA .² The latter complex contains a water molecule between the two Re(I) atoms. In contrast, $[\text{Re}(\text{CO})_3]_2\text{TPP}$ and $[\text{Re}(\text{CO})_3]_2\text{TAA}$ contain rhenium atoms above and below the donor plane but off the S_2 axis normal to the plane. X-Ray diffraction analysis have shown them to be centrosymmetric, having two metal atoms bound to three adjacent nitrogen atoms, one above and one below the plane of the macrocyclic ligand.^{2,15} The Re-Re separation is 3.345 \AA for the $[\text{Re}(\text{CO})_3]_2\text{TAA}$ complex² and 3.126 \AA for $[\text{Re}(\text{CO})_3]_2\text{TPP}$.¹⁵ These distances are too long for a formal bond but still allow some metal-metal interaction. [The $[\text{Re}(\text{CO})_3]_2\text{TAA}$ complex, unlike the only other TAA complex, NiTAA,¹⁶ studied by x-ray diffraction is non planar.³] The degree of twisting or distortion in the TAA framework of $[\text{Re}(\text{CO})_3]_2\text{TAA}$ is similar to that of the $[\text{Re}(\text{CO})_3]_2\text{TPP}$ compound. Also, the bond distances in the 2,4-pentanediiiminato side chain clearly show the nonlocalization of the macrocycle with alternating single and double bonds in the structure of $[\text{Re}(\text{CO})_3]_2\text{TAA}$. NiTAA on the other hand, shows a delocalized side chain. These factors lend structural uniqueness to $[\text{Re}(\text{CO})_3]_2\text{TAA}$ as a TAA compound.

It has been reported for octaethylporphyrin complexes,¹⁷ and verified with other metalloporphyrin complexes,¹⁸⁻²¹ that the potential difference between cation and anion radical formation, $|E_{1/2}(\text{ox}) - E_{1/2}(\text{red})|$, is equal to $2.25 \pm 0.15\text{V}$. With several exceptions, this value is independent of metal ion electron transfers and holds true for a number of different natural and synthetic porphyrins containing metals of +2, +3 or +4 oxidation state. Recently Gross and

Jordan²² showed that this potential difference decreased to 1.82V for $[\text{Re}(\text{CO})_3]_2$ TPP oxidation-reduction and that the change was due almost entirely to changes associated with the lowest unoccupied molecular orbital (LUMO). Our results with H_2TMTAA , $[\text{Re}(\text{CO})_3]_2\text{TMTAA}$ and $[\text{Rh}(\text{CO})_2]_2\text{TMTAA}$ also indicate a change in the LUMO but in this case, in a direction opposite to that observed for the porphyrin complexes. Surprisingly $[\text{Re}(\text{CO})_3]_2\text{TAA}$ behaves very similarly to most metalloporphyrins¹¹ in regards to the potential separation between the π radical reactions. The oxidation potential of 1.11 Volt in CH_2Cl_2 is almost identical to that for $[\text{Re}(\text{CO})_3]_2\text{TPP}$ and when combined with the $E_{1/2}(\text{red}) = -1.09$ in DMF yields an $|E_{1/2}(\text{ox}) - E_{1/2}(\text{red})| = 2.20\text{V}$.

Of special interest is a comparison of the $[\text{Rh}(\text{CO})_2]_2\text{TMTAA}$ with the $[\text{Re}(\text{CO})_3]_2\text{TMTAA}$. Both compounds are more easily oxidized than the uncomplexed ligand. However, the reversible half wave potential for oxidation of $[\text{Rh}(\text{CO})_2]_2\text{TMTAA}$ is anodic of $[\text{Re}(\text{CO})_3]_2\text{TMTAA}$ by 0.19V. This is opposite that predicted from the electronegativities of the two metal ions. For a series of M(II) to M(IV) metallocarboxyethylporphyrins, the difficulty of porphyrin ring oxidation is directly proportional to the negative charge on the porphyrin ligand, and for a given metal oxidation state, half wave potentials for cation radical formation can be linearly related to the electronegativities of the central metal.¹⁷ Since the trend in potentials is reversed for the tricarbonyl rhenium (I) and dicarbonyl rhodium (I), this would imply that between the two complexes differences in metal-metal interaction exist and that the interactions of the metal orbitals with those of the macrocyclic ring differ. Electrochemical studies of other mono and dimetallic complexes of H_2TMTAA are now underway to see if similar trends exist between oxidation potentials and degree of metal-metal interaction.

Acknowledgement.

Support from the Robert A. Welch Foundation (K.M.K.) and the Office of Naval Research (M.T.) is gratefully acknowledged.

References

1. (a) G.C. Gordon, P.W. DeHaven, M.C. Weiss and V.L. Goedken, J. Am. Chem. Soc., **100**, 1003(1978)
(b) V.L. Goedken, Personal communication.
2. M. Tsutsui, R.L. Bobsein and R. Pettersen, submitted for publication.
3. M. Tsutsui, R.L. Bobsein, R. Pettersen and R. Haaker, in press.
4. V.L. Goedken, J.J. Pluth, S.-M. Peng, and B. Bursten, J. Am. Chem. Soc., **98**, 8014(1976).
5. (a) H. Alt, H. Binder, and G. Sandstede, J. Catal., **28**, 8(1973).
(b) G.C. Demitras and E.L. Muetterties, J. Am. Chem. Soc., **99**, 2796(1977).
6. (a) V.L. Goedken, J. Molin-Case and Y-A. Whang, J. Chem. Soc., Chem. Commun., 337 (1973).
(b) V.L. Goedken and Y-A. Park., J. Chem. Soc., Chem. Commun., 214 (1975);
(c) D. R. Neves and J.C. Dabrowiak, Inorg. Chem., **15**, 129 (1976);
(d) M.C. Weiss and V.L. Goedken, J. Am. Chem. Soc., **98**, 3389 (1976).
7. (a) V.L. Goedken, Y-A. Park, S.-M. Peng and J.M. Norris, J. Am. Chem. Soc., **96**, 7693(1974); V.L. Goedken, S.-M. Peng, J.M. Norris, and Y-A. Park, J. Am. Chem. Soc., **98**, 8007(1976).
8. (a) W.H. Woodruff, R.W. Pastor and J.C. Dabrowiak, J. Am. Chem. Soc., **98**, 7999(1976); (b) L.A. Nafie, R.W. Pastor, J.C. Dabrowiak and W.H. Woodruff., J. Am. Chem. Soc., **98**, 8007(1976).
9. F.C. McElroy and J.C. Dabrowiak, J. Am. Chem. Soc., **98**, 7112(1976).
10. F.C. McElroy, J.C. Dabrowiak and D.J. Macero, Inorg. Chem., **16**, 947(1977).
11. J. Fuhrhop, "Porphyrins and Metalloporphyrins," K.M. Smith, Ed., Elsevier, New York, N.Y., 1975
12. R.S. Nicholson and I. Shain, Anal. Chem., **36**, 706(1964).
13. (a) M.C. Weiss, B. Bursten, S.-M. Peng, and V.L. Goedken, J. Chem. Soc., 8021(1976).
(b) M.C. Weiss and V.L. Goedken, J. Chem. Soc., Chem. Commun., 531(1976).
14. M. Tsutsui, R.L. Bobsein, G. Cash, and R. Pettersen, submitted for publication.
15. (a) D. Ostfeld, M. Tsutsui, C.P. Hrung and D.C. Conway, J. Am. Chem. Soc., **93**, 2548(1971);
(b) M. Tsutsui, D. Ostfeld and L.M. Hoffman, J. Am. Chem. Soc., **93**, 1820(1971)
(c) M. Tsutsui, C.P. Hrung, D. Ostfeld, T.S. Srivastava, D.L. Cullen, and E.F. Meyer, Jr., J. Am. Chem. Soc., **97**, 3952(1975).
16. M.C. Weiss, G. Gordon, and V.L. Goedken, Inorg. Chem., **16**, 305(1977).

17. J.-H. Fuhrhop, K.M. Kadish and D.G. Davis, J. Am. Chem. Soc., 95, 5140 (1973).
18. M.F. Jamin and R.T. Iwamoto, Inorg. Chem. Acta., 27, 135(1978).
19. K.M. Kadish and M.M. Morrison, Bioinorg. Chem., 7, 107(1977).
20. P. Fournari, R. Guillard, M. Fontesse, J.M. Latour and J.C. Marchon, J. Organomet. Chem., 110, 205 (1976).
21. K.M. Kadish, J.S. Cheng, I.A. Cohen and D. Summerville, ACS Symposium Series, 38, 65(1977).
22. A. Louati, M. Gross and J. Jordan, J. Electroanal. Chem., 86, 301(1978).

Table I. Half Wave Potentials of Cation and Anion Radical Formation in DMF

Compound	$E_{1/2}(\text{ox})$ (V vs. SLCE)	$E_{1/2}(\text{red})$ (V vs. SLCE)	$ E_{1/2}(\text{ox}) - E_{1/2}(\text{red}) $ (V)
H_2TPP	1.25	-0.90	2.15
H_2TAA	0.85 ^a	-1.80	2.60
H_2TMTAA	0.96 ^a	>-2.20	>3.16
$[\text{Re}(\text{CO})_3]_2\text{TPP}^b$	1.16	-0.66	1.82
$[\text{Re}(\text{CO})_3]_2\text{TAA}$	1.28 ^a	-1.09	2.37
$[\text{Re}(\text{CO})_3]_2\text{TMTAA}$	0.80	>-2.20	>3.00
$[\text{Rh}(\text{CO})_2]_2\text{TMTAA}$	0.61	>-2.20	>2.81

a. Anodic peak potential measured at a scan rate of 1.0 V/s


b. Taken from reference 22 where potentials were obtained in DMF containing 0.1M TBAP and are reported vs. SCE

Table II. Half Wave Potentials for Oxidation of Several Dirhenium and Dirhodium Carbonyl Complexes in CH_2Cl_2


Compound	$E_{1/2}$ (volt vs. SLCE)
$[\text{Rh}(\text{CO})_2]_2$ TMTAA	0.50
$[\text{Re}(\text{CO})_3]_2$ TMTAA	0.69
$[\text{Re}(\text{CO})_3]_2$ TAA	1.11
$[\text{Re}(\text{CO})_3]_2$ TPP	1.01

LIST OF FIGURES


1. Schematic Pictures of $[\text{Re}(\text{CO})_3]_2\text{TAA}$ (I), $[\text{Rh}(\text{CO})_2]_2\text{TMTAA}$ (II), $[\text{Re}(\text{CO})_3]_2\text{TPP}$ (III), and $[\text{Re}(\text{CO})_3]_2\text{TMTAA}$ (IV).
2. Cyclic voltammogram (solid line) and differential pulse polarogram (dashed line) showing cation and anion radical formation reactions of $1.8 \times 10^{-3}\text{M}$ H_2TAA in DMF, 0.1M TBAT. Scan rate = 0.2V/sec for cyclic voltammogram, 0.01 V/sec for differential pulse polarogram.




I



II



III



IV

CURRENT

